

Kaiser Permanente Research Brief


Aging

This brief summarizes the contributions of Kaiser Permanente Research since 2007 on the topic of aging. Although this topic encompasses a wide variety of health issues, this brief will focus on geriatric syndromes (for example, frailty, falls), cognitive illnesses, the management of multiple chronic conditions, and end-of-life care.

For much of the past century, average life expectancy in the United States has increased dramatically. That, combined with the aging of the baby boom generation, accounts for a growing number of older Americans, who now make up an unprecedented percentage of the population. Today, there are 35 million people in the United States age 65 and older, and by 2030, these older adults are expected to comprise 20% of the U.S. population. Additionally, there are 4 million people age 85 and older in the United States, who represent the fastest-growing segment of the population.¹ Of Americans age 65 and older, 2 out of every 3 have multiple chronic health problems.²

The health of older adults is complicated by common, multifactorial health problems known as geriatric syndromes. Frailty — a condition defined by fatigue and decreased strength, mobility, and activity — affects approximately 15% of older adults, with higher prevalence among women, members of racial and ethnic minorities, and lower-income individuals.³ According to the Centers for Disease Control and Prevention, 29 million American adults age 65 and older fall each year; these accidents are implicated in 30,000 deaths.⁴ Roughly half of older adults also suffer from urinary or bowel incontinence.⁵ Moreover, many adults experience declines in cognitive ability as they age; for many, these changes progress to dementia, a condition characterized by personality changes and impairments in memory and reasoning ability that limit their ability to live independently. The most common form of dementia, Alzheimer's disease, affects 1 in 9 Americans over the age of 65 and is associated with death and disability, significant burdens for paid and unpaid caregivers, and nearly \$345 billion in annual health care costs.⁶ Finally, the last months of life are frequently characterized by high medical costs⁷ and the challenge of balancing appropriate treatment and the patient's wishes. Although most patients with terminal illness prefer to die at home, recent data indicate that this only happens in 30% of cases.⁸

Aging is an active area of study for Kaiser Permanente Research. Scientists across the organization have used our rich, comprehensive, longitudinal data to advance knowledge in the areas of understanding risk, improving patient outcomes, and translating research findings into policy and practice. We have published more than 2,700 articles related to aging since 2007;⁹ together, these articles have been cited more than



This brief summarizes a selection of the publications contained within the Kaiser Permanente Publications Library, which indexes journal articles and other publications authored by individuals affiliated with Kaiser Permanente. The work described in this brief originated from across Kaiser Permanente's 8 regions and was supported by a wide range of funding sources including internal research support as well as both governmental and nongovernmental extramural funding.

106,000 times. These articles are the product of observational studies, randomized controlled trials, meta-analyses, and other studies led by Kaiser Permanente scientists. Our unique environment — a fully integrated care and coverage model in which our research scientists, clinicians, medical groups and health plan leaders collaborate — enables us to contribute generalizable knowledge on aging, and many other research topics.

Understanding risk

What are the unique health risks associated with aging?

Kaiser Permanente researchers have conducted extensive research on the unique health risks faced by older patients. Common illnesses such as osteoporosis,^{10, 11} chronic obstructive pulmonary disease,^{12, 13} and congestive heart failure^{14, 15} are more severe and have poorer prognoses at later ages. The challenge of managing these conditions is often increased in older patients, for whom the benefits of many medications must be weighed against heightened risks,¹⁶⁻²¹ and who are often underrepresented in clinical trials.²² For example, many widely prescribed medications are associated with heightened risks for falls,²³⁻²⁸ as well as declines in cognition²⁹⁻³¹ and functional status³⁰ in older patients.

One area of significant work at Kaiser Permanente concerns the challenges of managing multiple chronic conditions, particularly with respect to encouraging adherence to multiple medications among patients with cognitive impairment,³² and minimizing the burden of inappropriate care.³³ Use of multiple medications, known as polypharmacy, is common in patients with multiple chronic conditions, and is also associated with an increased risk of falls²⁶ and other adverse outcomes.^{34, 35} Sources of these risks include drug interactions³⁶⁻³⁸ and inadequate communication among multiple prescribing clinicians.³⁹ Polypharmacy and use of high-risk medications are also frequent causes of hospital-acquired delirium,⁴⁰ and many Kaiser Permanente medical centers have implemented early detection programs for delirium.⁴¹ Finally, treatment of multiple health problems often requires frequent medical imaging, which may lead to distress associated with incidental detection of new lesions.⁴²

Geriatric syndromes such as frailty, falls, and declines in functional ability are a focus of research at Kaiser Permanente. Studies conducted by our scientists have found that vitamin D deficiency,^{43, 44} disturbed sleep patterns,⁴⁵ vision impairment,⁴⁶ kidney disease,⁴⁷ low blood pressure,⁴⁸ blood lipid characteristics,⁴⁹ hypoglycemia associated with overuse of antidiabetic medications,^{50, 51} and metabolic illnesses^{52, 53} are associated with increased risks of falls and frailty. Functional status limitations, including disability and impaired mobility, are linked to disrupted sleep behaviors,⁵⁴ other geriatric syndromes,⁵⁵⁻⁵⁷ and increased risks of falls,⁵⁸⁻⁶⁰ cognitive impairment,⁶¹ hospitalization,⁶² bleeding risk,⁶³ and mortality.⁶⁴ Our scientists participated in a large, community-based study that found that lower walking speed and decreased physical activity were associated with chronic disability, long-term nursing home stays, and an increased risk of

Food insecurity

Nearly 6% of elderly Kaiser Permanente members in Colorado said that they did not always have enough money to buy food.¹⁰²



Less likely to experience food insecurity

- Age 85 and older
- Living at home with a friend or relative
- Heavy alcohol use
- Recent stay in a skilled nursing facility
- Someone to call for help

More likely to experience food insecurity

- Black Americans
- Medicaid or Special Needs Plan Insurance
- Fair or poor quality of life
- Dental or mouth problems
- Financial problems
- Zero daily serving of fruits or vegetables

death.⁶⁵ We have also explored various risk factors associated with age-related cognitive decline and dementia, including race or ethnicity,⁶⁶⁻⁶⁸ obesity,^{69, 70} physical inactivity,^{71, 72} poor pulmonary function,⁷³ sleep disturbances,^{74, 75} social isolation,⁷⁶ life stress,⁶⁶ brain characteristics,⁷⁷⁻⁷⁹ genetic factors,⁸⁰⁻⁸² environmental factors,⁸³⁻⁸⁵ and illnesses such as hypertension,^{86, 87} diabetes,⁸⁸⁻⁹² kidney disease,⁹³ and psychiatric disorders.⁹⁴⁻⁹⁷ Our scientists contributed to the Lancet Commission's life-course model for reducing dementia risks, which emphasizes early interventions to target modifiable risk factors.^{98, 99} This model was based in part on findings from the long-running Adult Changes in Thought study, conducted by Kaiser Permanente researchers in Washington.¹⁰⁰

Which subgroups of older adults are at particularly high risk for health problems?

Social needs are one domain of risk factors that impact older adults. Kaiser Permanente scientists have studied social determinants of health in older people.¹⁰¹ A survey of Kaiser Permanente members in Colorado age 65 and older found that approximately 6% did not have enough money for food. Black members and members with Medicaid insurance, lower self-reported quality of life, dental problems, poor diet, or inadequate social support were at particularly high risk.¹⁰² For older people, food insecurity is associated with poor diabetes control and higher rates of falls, hospitalizations, emergency department visits, and deaths.¹⁰³⁻¹⁰⁵ Financial constraints and poverty are also linked to poor health outcomes in older people, according to research conducted by our scientists.¹⁰⁶⁻¹⁰⁹ Kaiser Permanente studies also show that social isolation is associated with a lower quality of life in older people¹¹⁰ and with the development of cognitive illness⁷⁶ and cardiovascular disease.¹¹¹ These challenges have become particularly acute in recent years, as older people have faced unprecedented isolation and limitations on physical activity during the COVID-19 pandemic.¹¹² Our scientists have evaluated various methods for proactively screening our members for unmet social needs,¹¹³ and have found greater health care utilization in members with greater unmet needs.^{113, 114} Our researchers have also found that high-risk opioid prescribing in older patients is a common problem^{115, 116} and is associated with greater risks of fractures¹¹⁷ and overdoses.¹¹⁸

Improving Patient Outcomes

What prevention or early intervention strategies are effective in mitigating the health risks of aging?

Kaiser Permanente scientists have studied various programs for primary prevention among older members. One component of these efforts includes the promotion of general wellness. Kaiser Permanente research has supported the implementation of exercise and physical fitness programs, such as Silver Sneakers and the use of wearable fitness trackers, among older members.^{119, 120} Exercise programs in older patients are effective in preventing falls,¹²¹⁻¹²⁵ and may also reduce the risk of dementia^{72, 126} and reduce health care costs.¹²⁷ Our scientists have incorporated social supports into programs for older patients to limit the impact of social isolation,^{128, 129} and they have studied programs for early identification of depression.¹³⁰ The COVID-19 pandemic led to a rapid shift toward digital health tools, and our scientists have discussed the needs and challenges that older patients may experience in using these technologies effectively.¹³¹⁻¹³³ Recently, Kaiser Permanente researchers were involved in the evaluation of a nurse case management program for patients with Parkinson's disease.^{134, 135}

Researchers at Kaiser Permanente have explored efforts to expand advance care planning to ensure that patient preferences are honored at the end of life. A study of the Advanced Steps program, which allows patients to discuss their end-of-life priorities with a health care agent and a planning facilitator, found a strong association with high rates of preference-concordant care, and demonstrated that preferences may


evolve over time.¹³⁶ Other Kaiser Permanente scientists have studied the effect of incorporating advance care planning prompts into patients' electronic medical records.¹³⁷ Our researchers have explored ways to identify unrecognized dementia using medical record data,¹³⁸⁻¹⁴⁰ and have developed various methods for predicting the risk of future dementia^{88, 141-143} and other risks in patients with dementia.¹⁴⁴ A randomized trial is currently underway to evaluate whether advance care planning can increase the safety of glucose-lowering medications in older Kaiser members with type 2 diabetes.¹⁴⁵

What are the key factors in effective care for older adults?

The effectiveness and safety of medical care for older patients are significant areas of research at Kaiser Permanente. One component of effective care involves discontinuing or reducing the intensity of care for older members, as appropriate. For example, Kaiser Permanente research demonstrated that preventive breast,¹⁴⁶⁻¹⁴⁸ cervical,¹⁴⁹ and colorectal¹⁵⁰ cancer screenings have limited benefits in older patients. These findings have led to recommendations to curtail routine screenings at later ages to protect these patients from unnecessary tests and risks.

Our scientists have also studied medication safety efforts aimed at deprescribing treatments¹⁵¹⁻¹⁵⁵ that are associated with greater risks at older ages, including antidiabetic drugs,^{18, 156} statins,¹⁵⁷ anticoagulants,¹⁵⁸ and sedatives,^{159, 160} as well as the implementation of deprescribing efforts in hospice settings.¹⁶¹ The quality of communication between physicians and patients is critical for successful deprescribing.¹⁶²⁻¹⁶⁶ Kaiser Permanente researchers have also studied programs for reconciliation of medications among multiple clinicians. These programs ensure that older patients do not receive duplicative treatments.¹⁶⁷⁻¹⁷⁰ Our research has emphasized the importance of shared decision-making and goal setting in the care of these patients. Kaiser Permanente scientists have also explored barriers to use of nonpharmacologic pain treatments in older members, in lieu of prescription opioids.¹⁷¹

Finally, Kaiser Permanente researchers have studied the use of advance health care directives and other plans in end-of-life care. Our scientists have also studied the role of caregivers and surrogate decision-makers in the care of older patients with dementia.¹⁷² Studies conducted among our members in California have found that care delivered at the end of life is nearly always concordant with the preferences patients have expressed in their Physician Orders for Life-Sustaining Treatment forms. This research has also drawn attention to how preferences may change over time.^{136, 173} While advance directives are consistently associated with less frequent delivery of intensive care services and a lower likelihood of dying in a hospital,^{174, 175} these directives are often not completed. Kaiser Permanente scientists have also explored virtual methods for training family caregivers in the management of psychological and behavioral symptoms of dementia.¹⁷⁶



Translating Research Findings Into Policy and Practice

Kaiser Permanente is a learning health care system that works to systematically use research to inform and improve practice. Our research, clinical, and operational partners have tested a range of interventions to reduce risks and improve outcomes for older patients.¹⁷⁷ Scientists in Kaiser Permanente have studied the use of electronic patient portals to support both patients and caregivers at the end of life.¹⁷⁸⁻¹⁸¹ Our researchers were heavily involved in the development of Primary Care Plus, a program combining palliative care with pharmacy, social, and behavioral health services to holistically care for older patients with complex needs.¹⁸² This program was successfully implemented in Colorado¹⁸³ and is being expanded to other geographic areas within Kaiser Permanente.¹⁸⁴ Our scientists also led the Family Care Study, a randomized study that evaluated a preparedness, skill, enrichment, and predictability intervention for family caregivers of frail older adults.^{185, 186}

Our researchers also led the development and implementation of a nurse-led program for increasing palliative care engagement among patients with non-small cell lung cancer,^{187, 188} pharmacist-led medication deprescribing efforts,^{160, 167, 189, 190} closer integration of medical and dental care,¹⁹¹ and intensive care unit-based assessments of delirium.⁴¹ Kaiser Permanente researchers are also developing innovative methods for studying health problems in older patients. These include artificial intelligence models that have identified subgroups of older patients with consistent clusters of multiple chronic health problems¹⁹² and extensive survey studies of barriers to patient self-care.¹⁰⁷ Our scientists have also developed an algorithm for predicting hypoglycemic events among older patients with type 2 diabetes; this tool is now integrated into the electronic health record system in Northern California.¹⁹³ Finally, Kaiser Permanente researchers in Southern California developed an index that uses electronic medical records to improve the identification of older patients with chronic kidney disease who have health conditions that complicate the treatment of their kidney disease.¹⁹⁴

Kaiser Permanente research contributes to policy and practice change within our care delivery system, and has also advanced the national understanding of aging. Kaiser Permanente's research on aging since 2007 has been cited 620 times within recent consensus statements and clinical practice guidelines published by a wide range of entities, including the American Geriatrics Society.¹⁹⁵ In addition, Kaiser Permanente researchers and clinician scientists have directly contributed as authors of guidelines created for the U.S. Preventive Services Task Force,^{121, 122, 130, 196, 197} as well as a statement on palliative care from the American Thoracic Society and numerous partner organizations.¹⁹⁸ Our scientists have shown leadership in the broader field of research on aging. Their work includes extensive research on epidemiology, risk factors for development of dementia,^{66, 67, 69, 88, 96, 141, 142, 199-201} and health care in older patients with multiple chronic conditions.^{39, 107, 202-205} Kaiser Permanente researchers also have leading roles in the U.S. Deprescribing Research Network, a National Institute on Aging initiative aimed at producing and disseminating high-quality

Insights from Kaiser Permanente Research on Advance Care Planning



- Advance care plans have been associated with less frequent hospitalization, fewer in-hospital deaths, and greater use of hospice services.^{174;175;182}
- Embedding a Care Directives tab into the electronic health record, or conducting a consultation with trained facilitators, can increase documentation of patients' preferences for end-of-life care.^{136;137;173}

research regarding medication deprescribing in American adults.^{201, 206} Our scientists are involved in the national Creating Age-Friendly Health Systems initiative,²⁰⁷ and in an international effort to develop guidance for organizing multidisciplinary teams caring for patients with Parkinson's disease.²⁰⁸ A Kaiser Permanente scientist was also a lead author on a report on dementia caregiving published by the National Academies of Sciences, Engineering, and Medicine.²⁰⁹

Kaiser Permanente's 180 research scientists and 1,640 support staff members are based at 9 research centers. There are currently 2,160 studies underway, including clinical trials. Since 2007 our research scientists have published more than 23,000 articles in peer-reviewed journals. Kaiser Permanente currently serves approximately 12.5 million members in 8 states and the District of Columbia.

This brief was written by Nicholas P. Emptage, Anna C. Davis, and Elizabeth A. McGlynn. It is available online from about.kaiserpermanente.org/our-story/health-research/research-briefs. The authors wish to thank the following researchers for their contributions to the development of this brief: Richard W. Grant, Sascha Dublin, and Elizabeth A. Bayliss.

References

1. National Institute on Aging. Introduction. 2020; <https://www.nia.nih.gov/living-long-well-21st-century-strategic-directions-research-aging/introduction>. Accessed February 7, 2020.
2. Boersma P, Black LI, Ward BW. Prevalence of Multiple Chronic Conditions Among US Adults, 2018. *Preventing chronic disease*. 2020;17:E106.
3. Bandeen-Roche K, Seplaki CL, Huang J, et al. Frailty in Older Adults: A Nationally Representative Profile in the United States. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2015;70(11):1427-1434.
4. Bergen G, Stevens MR, Burns ER. Falls and Fall Injuries Among Adults Aged >=65 Years - United States, 2014. *MMWR Morbidity and mortality weekly report*. 2016;65(37):993-998.
5. Gorina Y, Schappert S, Bercovitz A, et al. *Prevalence of incontinence among older Americans*. National Center for Health Statistics;2014.
6. Alzheimer's Association. 2023 Alzheimer's disease facts and figures. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2023;19(4):1598-1695.
7. Riley GF, Lubitz JD. Long-term trends in Medicare payments in the last year of life. *Health services research*. 2010;45(2):565-576.
8. Cross SH, Warraich HJ. Changes in the Place of Death in the United States. *The New England journal of medicine*. 2019;381(24):2369-2370.
9. KPPL Search, conducted on November 9, 2023: ((Msubject:"dementia") OR (Msubject:"geriatric assessment") OR (Msubject:"geriatric psychiatry") OR (Msubject:"geriatric nursing") OR (Msubject:"geriatric dentistry") OR (Msubject:"health services for the aged") OR (Msubject:"palliative care") OR (Msubject:"hospice care") OR (Msubject:"advance care planning") OR (Msubject:"resuscitation orders") OR (Msubject:"aged") OR (Msubject:"frail elderly") OR (title:"dementia") OR (title:"Alzheimer**") OR (title:"Parkinson**") OR (title:"palliat**") OR (title:"hospice") OR (title:"elderly") OR (title:"oldest") OR (title:"geriatric")) NOT ((Msubject:"infant") OR (Msubject:"adolescent") OR (Msubject:"child") OR (Msubject:"middle aged") OR (Msubject:"young adult") OR (title:"pediatric**") OR (title:"paediatric**") OR (title:"newborn**") OR (title:"infan**") OR (title:"child**") OR (title:"boy**") OR (title:"girl**") OR (title:"teen**") OR (title:"adolescen**") OR (title:"young adult**") OR (title:"middle aged") OR (title:"middle-aged") OR (abstract:"pediatric**") OR (abstract:"paediatric**") OR (abstract:"newborn**") OR (abstract:"infan**") OR (abstract:"child**") OR (abstract:"boy**") OR (abstract:"girl**") OR (abstract:"teen**") OR (abstract:"adolescen**") OR (abstract:"young adult**") OR (abstract:"middle aged") OR (abstract:"middle-aged")) AND publicationtype:"Journal Article". Dates 2007 to 2023.

10. Mackey DC, Lui LY, Cawthon PM, et al. High-trauma fractures and low bone mineral density in older women and men. *JAMA*. 2007;298(20):2381-2388.
11. Cauley JA, Hochberg MC, Lui LY, et al. Long-term risk of incident vertebral fractures. *JAMA*. 2007;298(23):2761-2767.
12. Mannino DM, Sonia Buist A, Vollmer WM. Chronic obstructive pulmonary disease in the older adult: what defines abnormal lung function? *Thorax*. 2007;62(3):237-241.
13. Jackson ML, Walker R, Lee S, Larson E, Dublin S. Predicting 2-Year Risk of Developing Pneumonia in Older Adults without Dementia. *Journal of the American Geriatrics Society*. 2016;64(7):1439-1447.
14. Tisminetzky M, Gurwitz JH, Fan D, et al. Noncardiac-Related Morbidity, Mobility Limitation, and Outcomes in Older Adults With Heart Failure. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2020;75(10):1981-1988.
15. Tisminetzky M, Gurwitz JH, Fan D, et al. Multimorbidity Burden and Adverse Outcomes in a Community-Based Cohort of Adults with Heart Failure. *Journal of the American Geriatrics Society*. 2018;66(12):2305-2313.
16. McGrath ER, Go AS, Chang Y, et al. Use of Oral Anticoagulant Therapy in Older Adults with Atrial Fibrillation After Acute Ischemic Stroke. *Journal of the American Geriatrics Society*. 2017;65(2):241-248.
17. Caffiero N, Delate T, Ehizuelen MD, Vogel K. Effectiveness of a Clinical Pharmacist Medication Therapy Management Program in Discontinuation of Drugs to Avoid in the Elderly. *Journal of managed care & specialty pharmacy*. 2017;23(5):525-531.
18. Lee EA, Gibbs NE, Martin J, Ziel F, Polzin JK, Palmer-Toy D. Improving Care in Older Patients with Diabetes: A Focus on Glycemic Control. *The Permanente journal*. 2016;20(3):51-56.
19. McManus DD, Kiefe C, Lessard D, et al. Geriatric Conditions and Prescription of Vitamin K Antagonists vs. Direct Oral Anticoagulants Among Older Patients With Atrial Fibrillation: SAGE-AF. *Frontiers in cardiovascular medicine*. 2019;6:155.
20. Izano MA, Lo JC, Adams AL, et al. Bisphosphonate Treatment Beyond 5 Years and Hip Fracture Risk in Older Women. *JAMA network open*. 2020;3(12):e2025190.
21. Wang W, Lessard D, Kiefe CI, et al. Differential effect of anticoagulation according to cognitive function and frailty in older patients with atrial fibrillation. *Journal of the American Geriatrics Society*. 2023;71(2):394-403.
22. Lo CT, Niu F, Fredriks DA, Hui RL. Evaluation of the effectiveness and safety of Direct Oral Anticoagulants in Elderly Patients with Non-Valvular Atrial Fibrillation who are not Candidates for Warfarin in real world setting. *Journal of cardiovascular pharmacology*. 2022;79(1):e138-e143.
23. Spence MM, Shin PJ, Lee EA, Gibbs NE. Risk of injury associated with skeletal muscle relaxant use in older adults. *The Annals of pharmacotherapy*. 2013;47(7-8):993-998.
24. Shimbo D, Barrett Bowling C, Levitan EB, et al. Short-Term Risk of Serious Fall Injuries in Older Adults Initiating and Intensifying Treatment With Antihypertensive Medication. *Circulation Cardiovascular quality and outcomes*. 2016;9(3):222-229.
25. Huang ES, Karter AJ, Danielson KK, Warton EM, Ahmed AT. The association between the number of prescription medications and incident falls in a multi-ethnic population of adult type-2 diabetes patients: the diabetes and aging study. *Journal of general internal medicine*. 2010;25(2):141-146.
26. Green AR, Reifler LM, Bayliss EA, Weffald LA, Boyd CM. Drugs Contributing to Anticholinergic Burden and Risk of Fall or Fall-Related Injury among Older Adults with Mild Cognitive Impairment, Dementia and Multiple Chronic Conditions: A Retrospective Cohort Study. *Drugs & aging*. 2019;36(3):289-297.
27. Hart LA, Marcum ZA, Gray SL, Walker RL, Crane PK, Larson EB. The Association Between Central Nervous System-Active Medication Use and Fall-Related Injury in Community-Dwelling Older Adults with Dementia. *Pharmacotherapy*. 2019;39(5):530-543.
28. Moffet HH, Huang ES, Liu JY, et al. Severe hypoglycemia and falls in older adults with diabetes: The Diabetes & Aging Study. *Diabetes epidemiology and management*. 2023;12.
29. Koyama A, Steinman M, Ensrud K, Hillier TA, Yaffe K. Ten-Year Trajectory of Potentially Inappropriate Medications in Very Old Women: Importance of Cognitive Status. *Journal of the American Geriatrics Society*. 2013;61(2):258-263.

30. Koyama A, Steinman M, Ensrud K, Hillier TA, Yaffe K. Long-term Cognitive and Functional Effects of Potentially Inappropriate Medications in Older Women. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2014;69(4):423-429.
31. Green AR, Reifler LM, Boyd CM, Weffald LA, Bayliss EA. Medication Profiles of Patients with Cognitive Impairment and High Anticholinergic Burden. *Drugs & aging*. 2018;35(3):223-232.
32. Du Vall M, Weffald L, Delate T, Shetterly S, Bayliss EA. Clinical Factors Associated With Nonadherence to Chronic Medications in People With Cognitive Impairment. *The Senior care pharmacist*. 2022;37(5):191-199.
33. Ferris R, Blaum C, Kiwak E, et al. Perspectives of Patients, Clinicians, and Health System Leaders on Changes Needed to Improve the Health Care and Outcomes of Older Adults With Multiple Chronic Conditions. *Journal of aging and health*. 2018;30(5):778-799.
34. Kabue S, Liu V, Dyer W, Raebel M, Nichols G, Schmittiel J. Identifying Common Predictors of Multiple Adverse Outcomes Among Elderly Adults With Type-2 Diabetes. *Medical care*. 2019;57(9):702-709.
35. Karimi S, Dharia SP, Flora DS, Slattum PW. Anticholinergic burden: clinical implications for seniors and strategies for clinicians. *The Consultant pharmacist : the journal of the American Society of Consultant Pharmacists*. 2012;27(8):564-582.
36. Tisminetzky M, Bayliss EA, Magaziner JS, et al. Research Priorities to Advance the Health and Health Care of Older Adults with Multiple Chronic Conditions. *Journal of the American Geriatrics Society*. 2017;65(7):1549-1553.
37. Elston Lafata J, Simpkins J, Kaatz S, et al. What do medical records tell us about potentially harmful co-prescribing? *Joint Commission journal on quality and patient safety / Joint Commission Resources*. 2007;33(7):395-400.
38. Sanghai S, Wong C, Wang Z, et al. Rates of Potentially Inappropriate Dosing of Direct-Acting Oral Anticoagulants and Associations With Geriatric Conditions Among Older Patients With Atrial Fibrillation: The SAGE-AF Study. *Journal of the American Heart Association*. 2020;9(6):e014108.
39. Maciejewski ML, Hammill BG, Bayliss EA, et al. Prescriber Continuity and Disease Control of Older Adults. *Medical care*. 2017;55(4):405-410.
40. Lee EA, Gibbs NE, Fahey L, Whiffen TL. Making hospitals safer for older adults: updating quality metrics by understanding hospital-acquired delirium and its link to falls. *The Permanente journal*. 2013;17(4):32-36.
41. Adams CL, Scruth EA, Andrade C, et al. Implementing clinical practice guidelines for screening and detection of delirium in a 21-hospital system in northern california: real challenges in performance improvement. *Clinical nurse specialist CNS*. 2015;29(1):29-37.
42. Freiman MR, Clark JA, Slatore CG, et al. Patients' Knowledge, Beliefs, and Distress associated with Detection and Evaluation of Incidental Pulmonary Nodules for Cancer: Results from a Multicenter Survey. *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*. 2016;11(5):700-708.
43. Ensrud KE, Ewing SK, Fredman L, et al. Circulating 25-hydroxyvitamin D levels and frailty status in older women. *The Journal of clinical endocrinology and metabolism*. 2010;95(12):5266-5273.
44. LeBlanc ES, Zakher B, Daeges M, Pappas M, Chou R. Screening for Vitamin D Deficiency: A Systematic Review for the U.S. Preventive Services Task Force. *Annals of internal medicine*. 2015;162(2):109-122.
45. Stone KL, Ancoli-Israel S, Blackwell T, et al. Actigraphy-measured sleep characteristics and risk of falls in older women. *Archives of internal medicine*. 2008;168(16):1768-1775.
46. Coleman AL, Cummings SR, Yu F, et al. Binocular visual-field loss increases the risk of future falls in older white women. *Journal of the American Geriatrics Society*. 2007;55(3):357-364.
47. Bowling CB, Bromfield SG, Colantonio LD, et al. Association of Reduced eGFR and Albuminuria with Serious Fall Injuries among Older Adults. *Clinical journal of the American Society of Nephrology : CJASN*. 2016;11(7):1236-1243.
48. Sim JJ, Zhou H, Bhandari S, et al. Low Systolic Blood Pressure From Treatment and Association With Serious Falls/Syncope. *American journal of preventive medicine*. 2018;55(4):488-496.
49. Barzilay JI, Buzkova P, Kuller LH, et al. The Association of Lipids and Lipoproteins with Hip Fracture Risk: The Cardiovascular Health Study. *The American journal of medicine*. 2022;135(9):1101-1108.e1101.

50. Laiteerapong N, Karter AJ, Liu JY, et al. Correlates of quality of life in older adults with diabetes: the diabetes & aging study. *Diabetes care*. 2011;34(8):1749-1753.
51. Lipska KJ, Huang ES, Liu JY, et al. Glycemic control and diabetes complications across health status categories in older adults treated with insulin or insulin secretagogues: The Diabetes & Aging Study. *Journal of the American Geriatrics Society*. 2023.
52. Barzilay JI, Blaum C, Moore T, et al. Insulin resistance and inflammation as precursors of frailty: the Cardiovascular Health Study. *Archives of internal medicine*. 2007;167(7):635-641.
53. Barzilay JI, Stein PK. Association of the Metabolic Syndrome with Age-Related, Nonatherosclerotic, Chronic Medical Conditions. *Metabolic syndrome and related disorders*. 2011;9(5):327-335.
54. Spira AP, Covinsky K, Rebok GW, et al. Poor sleep quality and functional decline in older women. *Journal of the American Geriatrics Society*. 2012;60(6):1092-1098.
55. Rosso AL, Eaton CB, Wallace R, et al. Combined impact of geriatric syndromes and cardiometabolic diseases on measures of functional impairment. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2011;66(3):349-354.
56. Rosso AL, Eaton CB, Wallace R, et al. Geriatric Syndromes and Incident Disability in Older Women: Results from the Women's Health Initiative Observational Study. *Journal of the American Geriatrics Society*. 2013;61(3):371-379.
57. Kinnucan E, Molcjan MT, Wright DM, Switzer JA. A Prospective Look at the Link Between Frailty and Shoulder Function in Asymptomatic Elderly Individuals. *Geriatric orthopaedic surgery & rehabilitation*. 2018;9:2151459318777583.
58. Buchner DM, Rillamas-Sun E, Di C, et al. Accelerometer-Measured Moderate to Vigorous Physical Activity and Incidence Rates of Falls in Older Women. *Journal of the American Geriatrics Society*. 2017;65(11):2480-2487.
59. Bromfield SG, Ngameni CA, Colantonio LD, et al. Blood Pressure, Antihypertensive Polypharmacy, Frailty, and Risk for Serious Fall Injuries Among Older Treated Adults With Hypertension. *Hypertension (Dallas, Tex : 1979)*. 2017;70(2):259-266.
60. Ylitalo KR, Karvonen-Gutierrez CA, Sternfeld B, Pettee Gabriel K. Association of Physical Activity and Physical Functioning Phenotypes With Fall Risk Among Women. *Journal of aging and health*. 2021;33(5-6):409-417.
61. Farias ST, Lau K, Harvey D, Denny KG, Barba C, Mefford AN. Early Functional Limitations in Cognitively Normal Older Adults Predict Diagnostic Conversion to Mild Cognitive Impairment. *Journal of the American Geriatrics Society*. 2017;65(6):1152-1158.
62. Ensrud KE, Lui LY, Paudel ML, et al. Effects of Mobility and Cognition on Hospitalization and Inpatient Days in Women in Late Life. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2017;72(1):82-88.
63. Wang W, Saczynski JS, Lessard D, et al. Presence of Geriatric Conditions Is Prognostic of Major Bleeding in Older Patients with Atrial Fibrillation: a Cohort Study. *Journal of general internal medicine*. 2022.
64. Ensrud KE, Lui LY, Paudel ML, et al. Effects of Mobility and Cognition on Risk of Mortality in Women in Late Life: A Prospective Study. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2016;71(6):759-765.
65. Rothman MD, Leo-Summers L, Gill TM. Prognostic significance of potential frailty criteria. *Journal of the American Geriatrics Society*. 2008;56(12):2211-2116.
66. Gilsanz P, Quesenberry CP, Mayeda ER, Glymour MM, Farias ST, Whitmer RA. Stressors in Midlife and Risk of Dementia: The Role of Race and Education. *Alzheimer disease and associated disorders*. 2019;33(3):200-205.
67. Mayeda ER, Glymour MM, Quesenberry CP, Whitmer RA. Heterogeneity in 14-year Dementia Incidence Between Asian American Subgroups. *Alzheimer disease and associated disorders*. 2017;31(3):181-186.
68. Gilsanz P, Corrada MM, Kawas CH, et al. Incidence of dementia after age 90 in a multiracial cohort. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2019;15(4):497-505.
69. Whitmer RA. The epidemiology of adiposity and dementia. *Current Alzheimer research*. 2007;4(2):117-122.

70. Lee CM, Woodward M, Batty GD, et al. Association of anthropometry and weight change with risk of dementia and its major subtypes: A meta-analysis consisting 2.8 million adults with 57 294 cases of dementia. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2020;21(4):e12989.
71. Barnes DE, Blackwell T, Stone KL, et al. Cognition in older women: the importance of daytime movement. *Journal of the American Geriatrics Society*. 2008;56(9):1658-1664.
72. Barnes DE, Whitmer RA, Yaffe K. Physical activity and dementia: The need for prevention trials. *Exercise and sport sciences reviews*. 2007;35(1):24-29.
73. Gilsanz P, Mayeda ER, Flatt J, Glymour MM, Quesenberry CP, Whitmer RA. Early Midlife Pulmonary Function and Dementia Risk. *Alzheimer disease and associated disorders*. 2018;32(4):270-275.
74. Tranah GJ, Blackwell T, Stone KL, et al. Circadian activity rhythms and risk of incident dementia and mild cognitive impairment in older women. *Annals of neurology*. 2011;70(5):722-732.
75. Diem SJ, Blackwell TL, Stone KL, et al. Measures of Sleep-Wake Patterns and Risk of Mild Cognitive Impairment or Dementia in Older Women. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2016;24(3):248-258.
76. Crooks VC, Lubben J, Petitti DB, Little D, Chiu V. Social network, cognitive function, and dementia incidence among elderly women. *American journal of public health*. 2008;98(7):1221-1227.
77. Fessel J. Amyloid is essential but insufficient for Alzheimer causation: addition of subcellular cofactors is required for dementia. *International journal of geriatric psychiatry*. 2018;33(1):e14-e21.
78. Fessel WJ. Concordance of Several Subcellular Interactions Initiates Alzheimer's Dementia: Their Reversal Requires Combination Treatment. *American journal of Alzheimer's disease and other dementias*. 2017;32(3):166-181.
79. Nelson PT, Brayne C, Flanagan ME, et al. Frequency of LATE neuropathologic change across the spectrum of Alzheimer's disease neuropathology: combined data from 13 community-based or population-based autopsy cohorts. *Acta neuropathologica*. 2022;144(1):27-44.
80. Kunkle BW, Grenier-Boley B, Sims R, et al. Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates A β , tau, immunity and lipid processing. *Nature genetics*. 2019;51(3):414-430.
81. Reiman EM, Arboleda-Velasquez JF, Quiroz YT, et al. Exceptionally low likelihood of Alzheimer's dementia in APOE2 homozygotes from a 5,000-person neuropathological study. *Nature communications*. 2020;11(1):667.
82. Dumitrescu L, Mahoney ER, Mukherjee S, et al. Genetic variants and functional pathways associated with resilience to Alzheimer's disease. *Brain : a journal of neurology*. 2020;143(8):2561-2575.
83. Iaccarino L, La Joie R, Lesman-Segev OH, et al. Association Between Ambient Air Pollution and Amyloid Positron Emission Tomography Positivity in Older Adults With Cognitive Impairment. *JAMA neurology*. 2021;78(2):197-207.
84. Shaffer RM, Blanco MN, Li G, et al. Fine Particulate Matter and Dementia Incidence in the Adult Changes in Thought Study. *Environmental health perspectives*. 2021;129(8):87001.
85. Shaffer RM, Li G, Adar SD, et al. Fine Particulate Matter and Markers of Alzheimer's Disease Neuropathology at Autopsy in a Community-Based Cohort. *Journal of Alzheimer's disease : JAD*. 2021;79(4):1761-1773.
86. van Dalen JW, Brayne C, Crane PK, et al. Association of Systolic Blood Pressure With Dementia Risk and the Role of Age, U-Shaped Associations, and Mortality. *JAMA internal medicine*. 2022;182(2):142-152.
87. Abdulrahman H, van Dalen JW, den Brok M, Latimer CS, Larson EB, Richard E. Hypertension and Alzheimer's disease pathology at autopsy: A systematic review. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2022.
88. Exalto LG, Biessels GJ, Karter AJ, et al. Risk score for prediction of 10 year dementia risk in individuals with type 2 diabetes: a cohort study. *The lancet Diabetes & endocrinology*. 2013;1(3):183-190.
89. Exalto LG, Whitmer RA, Kappele LJ, Biessels GJ. An update on type 2 diabetes, vascular dementia and Alzheimer's disease. *Experimental gerontology*. 2012;47(11):858-864.
90. Mayeda ER, Whitmer RA, Yaffe K. Diabetes and Cognition. *Clinics in geriatric medicine*. 2015;31(1):101-115, ix.
91. Lacy ME, Gilsanz P, Karter AJ, Quesenberry CP, Pletcher MJ, Whitmer RA. Long-term Glycemic Control and Dementia Risk in Type 1 Diabetes. *Diabetes care*. 2018;41(11):2339-2345.

92. Moran C, Lacy ME, Whitmer RA, et al. Glycemic Control Over Multiple Decades and Dementia Risk in People With Type 2 Diabetes. *JAMA neurology*. 2023;80(6):597-604.
93. Barzilay JI, Fitzpatrick AL, Luchsinger J, et al. Albuminuria and dementia in the elderly: a community study. *American journal of kidney diseases : the official journal of the National Kidney Foundation*. 2008;52(2):216-226.
94. Colenda CC, Legault C, Rapp SR, et al. Psychiatric disorders and cognitive dysfunction among older, postmenopausal women: results from the Women's Health Initiative Memory Study. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2010;18(2):177-186.
95. Flatt JD, Gilsanz P, Quesenberry CP, Albers KB, Whitmer RA. Post-traumatic stress disorder and risk of dementia among members of a health care delivery system. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2018;14(1):28-34.
96. Gilsanz P, Schnaider Beerli M, Karter AJ, Quesenberry CP, Adams AS, Whitmer RA. Depression in type 1 diabetes and risk of dementia. *Aging & mental health*. 2019;23(7):880-886.
97. Lowe XR, Merchant M, R AW. Antipsychotic Medication Risk of Dementia and Death: A Propensity Matched Cohort Study. *Current drug research reviews*. 2022;14(2):139-147.
98. Livingston G, Sommerlad A, Orgeta V, et al. Dementia prevention, intervention, and care. *Lancet (London, England)*. 2017;390(10113):2673-2734.
99. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet (London, England)*. 2020;396(10248):413-446.
100. ACT study: Long-running study of aging examines changes in Kaiser Permanente patients over time [press release]. 2018.
101. Bayliss EA, Ellis JL, Powers JD, Gozansky W, Zeng C. Using Self-Reported Data to Segment Older Adult Populations with Complex Care Needs. *EGEMS (Washington, DC)*. 2019;7(1):12.
102. Steiner JF, Stenmark SH, Sterrett AT, et al. Food Insecurity in Older Adults in an Integrated Health Care System. *Journal of the American Geriatrics Society*. 2018;66(5):1017-1024.
103. Schroeder EB, Zeng C, Sterrett AT, Kimpo TK, Paolino AR, Steiner JF. The longitudinal relationship between food insecurity in older adults with diabetes and emergency department visits, hospitalizations, hemoglobin A1c, and medication adherence. *Journal of diabetes and its complications*. 2019;33(4):289-295.
104. Mosen DM, Banegas MP, Friedman N, Shuster E, Brooks N. Food Insecurity Associated with Self-Reported Falls Among Medicare Advantage Members. *Population health management*. 2019;22(6):536-539.
105. Nguyen HQ, Duan L, Lee JS, et al. Association of a Medicare Advantage Posthospitalization Home Meal Delivery Benefit With Rehospitalization and Death. *JAMA health forum*. 2023;4(6):e231678.
106. Fung V, Reed M, Price M, et al. Responses to Medicare Drug Costs among Near-Poor versus Subsidized Beneficiaries. *Health services research*. 2013;48(5):1653-1668.
107. Bayliss EA, Ellis JL, Steiner JF. Barriers to self-management and quality-of-life outcomes in seniors with multimorbidities. *Annals of family medicine*. 2007;5(5):395-402.
108. Gerber Y, Weston SA, Killian JM, Therneau TM, Jacobsen SJ, Roger VL. Neighborhood income and individual education: effect on survival after myocardial infarction. *Mayo Clinic proceedings*. 2008;83(6):663-669.
109. Lee DR, Santo EC, Lo JC, Ritterman Weintraub ML, Patton M, Gordon NP. Understanding functional and social risk characteristics of frail older adults: a cross-sectional survey study. *BMC family practice*. 2018;19(1):170.
110. Maxwell CJ, Kang J, Walker JD, et al. Sex differences in the relative contribution of social and clinical factors to the Health Utilities Index Mark 2 measure of health-related quality of life in older home care clients. *Health and quality of life outcomes*. 2009;7:80.
111. Golaszewski NM, LaCroix AZ, Godino JG, et al. Evaluation of Social Isolation, Loneliness, and Cardiovascular Disease Among Older Women in the US. *JAMA network open*. 2022;5(2):e2146461.
112. Greenwood-Hickman MA, Dahlquist J, Cooper J, et al. "They're Going to Zoom It": A Qualitative Investigation of Impacts and Coping Strategies During the COVID-19 Pandemic Among Older Adults. *Frontiers in public health*. 2021;9:679976.

113. Schickedanz A, Sharp A, Hu YR, et al. Impact of Social Needs Navigation on Utilization Among High Utilizers in a Large Integrated Health System: a Quasi-experimental Study. *Journal of general internal medicine*. 2019;34(11):2382-2389.
114. Rogers A, Hu YR, Schickedanz A, Gottlieb L, Sharp A. Understanding High-Utilizing Patients Based on Social Risk Profiles: a Latent Class Analysis Within an Integrated Health System. *Journal of general internal medicine*. 2020;35(7):2214-2216.
115. Raman SR, Bush C, Karmali RN, Greenblatt LH, Roberts AW, Skinner AC. Characteristics of New Opioid Use Among Medicare Beneficiaries: Identifying High-Risk Patterns. *Journal of managed care & specialty pharmacy*. 2019;25(9):966-972.
116. Gold LS, Hansen RN, Avins AL, et al. Associations of Early Opioid Use with Patient-Reported Outcomes and Healthcare Utilization Among Older Adults with Low Back Pain. *The Clinical journal of pain*. 2018;34(4):297-305.
117. Saunders KW, Dunn KM, Merrill JO, et al. Relationship of opioid use and dosage levels to fractures in older chronic pain patients. *Journal of general internal medicine*. 2010;25(4):310-315.
118. Peglow SL, Binswanger IA. Preventing Opioid Overdose in the Clinic and Hospital: Analgesia and Opioid Antagonists. *The Medical clinics of North America*. 2018;102(4):621-634.
119. Dunn AL, Buller DB, Dearing JW, et al. Adopting an Evidence-Based Lifestyle Physical Activity Program: Dissemination Study Design and Methods. *Translational behavioral medicine*. 2012;2(2):199-208.
120. Rosenberg D, Walker R, Greenwood-Hickman MA, et al. Device-assessed physical activity and sedentary behavior in a community-based cohort of older adults. *BMC public health*. 2020;20(1):1256.
121. Guirguis-Blake JM, Michael YL, Perdue LA, Coppola EL, Beil TL. Interventions to Prevent Falls in Older Adults: Updated Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA*. 2018;319(16):1705-1716.
122. Force USPST, Grossman DC, Curry SJ, et al. Interventions to Prevent Falls in Community-Dwelling Older Adults: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2018;319(16):1696-1704.
123. Paganini-Hill A, Greenia DE, Perry S, Sajjadi SA, Kawas CH, Corrada MM. Lower likelihood of falling at age 90+ is associated with daily exercise a quarter of a century earlier: The 90+ Study. *Age and ageing*. 2017;46(6):951-957.
124. Li F, Harmer P, Glasgow R, et al. Translation of an effective tai chi intervention into a community-based falls-prevention program. *American journal of public health*. 2008;98(7):1195-1198.
125. Greenwood-Hickman MA, Rosenberg DE, Phelan EA, Fitzpatrick AL. Participation in Older Adult Physical Activity Programs and Risk for Falls Requiring Medical Care, Washington State, 2005-2011. *Preventing chronic disease*. 2015;12:E90.
126. LaHue SC, Comella CL, Tanner CM. The best medicine? The influence of physical activity and inactivity on Parkinson's disease. *Movement disorders : official journal of the Movement Disorder Society*. 2016;31(10):1444-1454.
127. Nguyen HQ, Maciejewski ML, Gao S, Lin E, Williams B, Logerfo JP. Health care use and costs associated with use of a health club membership benefit in older adults with diabetes. *Diabetes care*. 2008;31(8):1562-1567.
128. Brumley R, Enguidanos S, Jamison P, et al. Increased satisfaction with care and lower costs: results of a randomized trial of in-home palliative care. *Journal of the American Geriatrics Society*. 2007;55(7):993-1000.
129. Yaffe K, Barnes DE, Rosenberg D, et al. Systematic Multi-Domain Alzheimer's Risk Reduction Trial (SMARRT): Study Protocol. *Journal of Alzheimer's disease : JAD*. 2019;70(S1):S207-S220.
130. O'Connor EA, Whitlock EP, Gaynes B, Beil TL. *Screening for Depression in Adults and Older Adults in Primary Care: An Updated Systematic Review*. Rockville, MD: Agency for Healthcare Research and Quality; 2009-12 2009.
131. Krishnaswami A, Beavers C, Dorsch MP, et al. Gerotechnology for Older Adults With Cardiovascular Diseases: JACC State-of-the-Art Review. *Journal of the American College of Cardiology*. 2020;76(22):2650-2670.
132. Steele Gray C, Chau E, Tahsin F, et al. Assessing the Implementation and Effectiveness of the Electronic Patient-Reported Outcome Tool for Older Adults With Complex Care Needs: Mixed Methods Study. *Journal of medical Internet research*. 2021;23(12):e29071.

133. McCloskey JK, Ellis JL, Uratsu CS, et al. Telehealth During COVID-19 for Adults with Multiple Chronic Conditions: Associations with Self-Reported Food Insecurity and with Physical Limitations. *Telemedicine journal and e-health : the official journal of the American Telemedicine Association*. 2023;29(10):1446-1454.
134. Connor KI, Siebens HC, Mittman BS, et al. Quality and extent of implementation of a nurse-led care management intervention: care coordination for health promotion and activities in Parkinson's disease (CHAPS). *BMC health services research*. 2020;20(1):732.
135. Connor KI, Siebens HC, Mittman BS, et al. Implementation fidelity of a nurse-led RCT-tested complex intervention, care coordination for health promotion and activities in Parkinson's disease (CHAPS) in meeting challenges in care management. *BMC neurology*. 2022;22(1):36.
136. Hopping-Winn J, Mullin J, March L, Caughey M, Stern M, Jarvie J. The Progression of End-of-Life Wishes and Concordance with End-of-Life Care. *Journal of palliative medicine*. 2018;21(4):541-545.
137. Turley M, Wang S, Meng D, Kanter M, Garrido T. Impact of a Care Directives Activity Tab in the Electronic Health Record on Documentation of Advance Care Planning. *The Permanente journal*. 2016;20(2):43-48.
138. Barnes DE, Zhou J, Walker RL, et al. Development and Validation of eRADAR: A Tool Using EHR Data to Detect Unrecognized Dementia. *Journal of the American Geriatrics Society*. 2020;68(1):103-111.
139. Harding BN, Floyd JS, Scherrer JF, et al. Methods to identify dementia in the electronic health record: Comparing cognitive test scores with dementia algorithms. *Healthcare (Amsterdam, Netherlands)*. 2020;8(2):100430.
140. Coley RY, Smith JJ, Karliner L, et al. External Validation of the eRADAR Risk Score for Detecting Undiagnosed Dementia in Two Real-World Healthcare Systems. *Journal of general internal medicine*. 2022.
141. Barnes DE, Covinsky KE, Whitmer RA, Kuller LH, Lopez OL, Yaffe K. Predicting risk of dementia in older adults: The late-life dementia risk index. *Neurology*. 2009;73(3):173-179.
142. Exalto LG, Quesenberry CP, Barnes D, Kivipelto M, Biessels GJ, Whitmer RA. Midlife risk score for the prediction of dementia four decades later. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2014;10(5):562-570.
143. Robertson K, Larson EB, Crane PK, et al. Using Varying Diagnostic Criteria to Examine Mild Cognitive Impairment Prevalence and Predict Dementia Incidence in a Community-Based Sample. *Journal of Alzheimer's disease : JAD*. 2019;68(4):1439-1451.
144. Suri JS, Maindarkar MA, Paul S, et al. Deep Learning Paradigm for Cardiovascular Disease/Stroke Risk Stratification in Parkinson's Disease Affected by COVID-19: A Narrative Review. *Diagnostics (Basel, Switzerland)*. 2022;12(7):1543.
145. [no authors]. Study Details | Prepare for Your Diabetes Care. 2022; <https://clinicaltrials.gov/study/NCT05263310>. Accessed December 5, 2023.
146. Demb J, Abraham L, Miglioretti DL, et al. Screening mammography outcomes: risk of breast cancer and mortality by comorbidity score and age. *Journal of the National Cancer Institute*. 2020;112(6):599-606.
147. Elmore JG, Ngo-Metzger Q. The Life and Death of Mammograms in Patients 75 Years and Older-To Screen or Not to Screen? *JAMA internal medicine*. 2020;180(6):843-844.
148. Schousboe JT, Sprague BL, Abraham L, et al. Cost-Effectiveness of Screening Mammography Beyond Age 75 Years : A Cost-Effectiveness Analysis. *Annals of internal medicine*. 2022;175(1):11-19.
149. Vesco KK, Whitlock EP, Eder M, Burda BU, Senger CA, Lutz K. Risk Factors and Other Epidemiologic Considerations for Cervical Cancer Screening: A Narrative Review for the U.S. Preventive Services Task Force. *Annals of internal medicine*. 2011;155(10):698-705, W216.
150. Wu BU. Selection of elderly patients for colorectal cancer screening. *JAMA internal medicine*. 2015;175(5):860-861.
151. Krishnaswami A, Steinman MA, Goyal P, et al. Deprescribing in Older Adults With Cardiovascular Disease. *Journal of the American College of Cardiology*. 2019;73(20):2584-2595.
152. Hahn EE, Munoz-Plaza CE, Lee EA, et al. Patient and Physician Perspectives of Deprescribing Potentially Inappropriate Medications in Older Adults with a History of Falls: a Qualitative Study. *Journal of general internal medicine*. 2021;36(10):3015-3022.
153. Bayliss EA, Albers K, Gleason K, et al. Recommendations for outcome measurement for deprescribing intervention studies. *Journal of the American Geriatrics Society*. 2022;70(9):2487-2497.

154. Bayliss EA, Shetterly SM, Drace ML, et al. Deprescribing Education vs Usual Care for Patients With Cognitive Impairment and Primary Care Clinicians: The OPTIMIZE Pragmatic Cluster Randomized Trial. *JAMA internal medicine*. 2022;182(5):534-542.
155. Lee EA, Wong CA, Barrio L, et al. An Approach to Deprescribe Antidepressants for Depression in Older Adults: Consensus, Multidisciplinary Practice Recommendations. *The Permanente journal*. 2023;27(2):1-8.
156. Weiner JZ, Gopalan A, Mishra P, et al. Use and Discontinuation of Insulin Treatment Among Adults Aged 75 to 79 Years With Type 2 Diabetes. *JAMA internal medicine*. 2019;179(12):1633-1641.
157. Gurwitz JH, Go AS, Fortmann SP. Statins for Primary Prevention in Older Adults: Uncertainty and the Need for More Evidence. *JAMA*. 2016;316(19):1971-1972.
158. Shah SJ, Singer DE, Fang MC, Reynolds K, Go AS, Eckman MH. Net Clinical Benefit of Oral Anticoagulation Among Older Adults With Atrial Fibrillation. *Circulation Cardiovascular quality and outcomes*. 2019;12(11):e006212.
159. Kuntz J, Kouch L, Christian D, Peterson PL, Gruss I. Barriers and Facilitators to the Deprescribing of Nonbenzodiazepine Sedative Medications Among Older Adults. *The Permanente journal*. 2018;22:17-157.
160. Kuntz JL, Kouch L, Christian D, Hu W, Peterson PL. Patient Education and Pharmacist Consultation Influence on Nonbenzodiazepine Sedative Medication Deprescribing Success for Older Adults. *The Permanente journal*. 2019;23:18-161.
161. Le V, Patel N, Nguyen Q, et al. Retrospective analysis of a pilot pharmacist-led hospice deprescribing program initiative. *Journal of the American Geriatrics Society*. 2021;69(5):1370-1376.
162. Goyal P, Anderson TS, Bernacki GM, et al. Physician Perspectives on Deprescribing Cardiovascular Medications for Older Adults. *Journal of the American Geriatrics Society*. 2020;68(1):78-86.
163. Reeve E, Wolff JL, Skehan M, Bayliss EA, Hilmer SN, Boyd CM. Assessment of Attitudes Toward Deprescribing in Older Medicare Beneficiaries in the United States. *JAMA internal medicine*. 2018;178(12):1673-1680.
164. Green AR, Boyd CM, Gleason KS, et al. Designing a Primary Care-Based Deprescribing Intervention for Patients with Dementia and Multiple Chronic Conditions: a Qualitative Study. *Journal of general internal medicine*. 2020;35(12):3556-3563.
165. Norton JD, Zeng C, Bayliss EA, et al. Ethical Aspects of Physician Decision-Making for Deprescribing Among Older Adults With Dementia. *JAMA network open*. 2023;6(10):e2336728.
166. Reeve E, Bayliss EA, Shetterly S, et al. Willingness of older people living with dementia and mild cognitive impairment and their caregivers to have medications deprescribed. *Age and ageing*. 2023;52(1).
167. Delate T, Chester EA, Stubbings TW, Barnes CA. Clinical outcomes of a home-based medication reconciliation program after discharge from a skilled nursing facility. *Pharmacotherapy*. 2008;28(4):444-452.
168. Fleisher J, Barbosa W, Sweeney MM, et al. Interdisciplinary Home Visits for Individuals with Advanced Parkinson's Disease and Related Disorders. *Journal of the American Geriatrics Society*. 2018;66(6):1226-1232.
169. Maciejewski ML, Hammill BG, Voils CI, et al. Prescriber continuity and medication availability in older adults with cardiometabolic conditions. *SAGE open medicine*. 2018;6:2050312118757388.
170. McNicholl IR, Gandhi M, Hare CB, Greene M, Pierluissi E. A Pharmacist-Led Program to Evaluate and Reduce Polypharmacy and Potentially Inappropriate Prescribing in Older, HIV-Positive Patients. *Pharmacotherapy*. 2017;37(12):1498-1506.
171. Karmali RN, Skinner AC, Trogdon JG, Weinberger M, George SZ, Hassmiller Lich K. The association between the supply of select nonpharmacologic providers for pain and use of nonpharmacologic pain management services and initial opioid prescribing patterns for Medicare beneficiaries with persistent musculoskeletal pain. *Health services research*. 2021;56(2):275-288.
172. Chodosh J, Mittman BS, Connor KI, et al. Caring for patients with dementia: how good is the quality of care? Results from three health systems. *Journal of the American Geriatrics Society*. 2007;55(8):1260-1268.
173. Turley M, Wang S, Meng D, Kanter MH, Garrido T. An information model for automated assessment of concordance between advance care preferences and care delivered near the end of life. *Journal of the American Medical Informatics Association : JAMIA*. 2016;23(e1):e118-124.

174. Jennings LA, Turner M, Keebler C, et al. The Effect of a Comprehensive Dementia Care Management Program on End-of-Life Care. *Journal of the American Geriatrics Society*. 2019;67(3):443-448.
175. Kurella Tamura M, Montez-Rath ME, Hall YN, Katz R, O'Hare AM. Advance Directives and End-of-Life Care among Nursing Home Residents Receiving Maintenance Dialysis. *Clinical journal of the American Society of Nephrology : CJASN*. 2017;12(3):435-442.
176. Ramirez M, Duran MC, Pabiniak CJ, et al. Family Caregiver Needs and Preferences for Virtual Training to Manage Behavioral and Psychological Symptoms of Dementia: Interview Study. *JMIR aging*. 2021;4(1):e24965.
177. Tuzzio L, Hanson LR, Reuben DB, et al. Transforming Dementia Care Through Pragmatic Clinical Trials Embedded in Learning Healthcare Systems. *Journal of the American Geriatrics Society*. 2020;68 Suppl 2:S43-S48.:S43-S48.
178. Dickman Portz J, Powers JD, Casillas A, et al. Characteristics of Patients and Proxy Caregivers Using Patient Portals in the Setting of Serious Illness and End of Life. *Journal of palliative medicine*. 2021;24(11):1697-1704.
179. Portz JD, Lum HD, Bull S, et al. Perceptions of Patient Portal Use for Advance Directive Documentation among Older Adults with Multiple Chronic Conditions. *Journal of social work in end-of-life & palliative care*. 2020;16(3):238-249.
180. Portz JD, Powers JD, Baldwin M, et al. Patient Portal Use Near the End-of-Life. *Journal of general internal medicine*. 2021.
181. Portz JD, Powers JD, Baldwin M, et al. How can patient portals support caregivers of older adults near the end-of-life? A mixed-methods study. *Journal of the American Geriatrics Society*. 2022;70(8):2431-2435.
182. Gozansky W. Is there a medical impact in addressing non-medical needs? A new approach to primary care involving complex cases says 'yes'. 2018; <https://permanente.org/medical-impact-addressing-non-medical-needs/>. Accessed July 23, 2020.
183. Kanter M, Courneya PT. Perspective on Publishing Quality Improvement Efforts. *The Permanente journal*. 2017;21.:17-140.
184. Kanter MH, Courneya PT. The Importance of Continual Learning in a Rapidly Changing Health Care Environment. *The Permanente journal*. 2018;22:18-071.
185. Hagen CM, Archbold PG, Miller LL, et al. How Tailoring Led to Variation in Care Issues, Dosage, and Outcomes: Part 1: Secondary Analysis of the PREP Trial for Frail Older Adults and Family Caregivers. *Research in gerontological nursing*. 2023;16(2):57-70.
186. Stewart BJ, Lyons KS, Hornbrook MC, et al. How Concurrent Services Obscured Detection of Intervention Benefits: Part 2: Secondary Analysis of the PREP Trial for Frail Older Adults and Family Caregivers. *Research in gerontological nursing*. 2023;16(2):71-83.
187. Nguyen HQ, Cuyegkeng T, Phung TO, et al. Integration of a Palliative Care Intervention into Community Practice for Lung Cancer: A Study Protocol and Lessons Learned with Implementation. *Journal of palliative medicine*. 2017;20(12):1327-1337.
188. Nguyen HQ, Ruel N, Macias M, et al. Translation and Evaluation of a Lung Cancer, Palliative Care Intervention for Community Practice. *Journal of pain and symptom management*. 2018;56(5):709-718.
189. Hui RL, Chang CC, Niu F, et al. Evaluation of a Pharmacist-Managed Antidiabetic Deprescribing Program in an Integrated Health Care System. *Journal of managed care & specialty pharmacy*. 2019;25(8):927-934.
190. Bayliss EA, Shetterly SM, Drace ML, et al. The OPTIMIZE patient- and family-centered, primary care-based deprescribing intervention for older adults with dementia or mild cognitive impairment and multiple chronic conditions: study protocol for a pragmatic cluster randomized controlled trial. *Trials*. 2020;21(1):542.
191. Mosen DM, Banegas MP, Dickerson JF, et al. Examining the association of medical-dental integration with closure of medical care gaps among the elderly population. *Journal of the American Dental Association (1939)*. 2021;152(4):302-308.
192. Grant RW, McCloskey J, Hatfield M, et al. Use of Latent Class Analysis and k-Means Clustering to Identify Complex Patient Profiles. *JAMA network open*. 2020;3(12):e2029068.

193. Karter AJ, Warton EM, Lipska KJ, et al. Development and Validation of a Tool to Identify Patients With Type 2 Diabetes at High Risk of Hypoglycemia-Related Emergency Department or Hospital Use. *JAMA internal medicine*. 2017;177(10):1461-1470.
194. Hall RK, Zhou H, Reynolds K, Harrison TN, Bowling CB. A Novel Approach to Developing a Discordance Index for Older Adults With Chronic Kidney Disease. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2020;75(3):522-528.
195. American Geriatrics Society 2012 Beers Criteria Update Expert Panel. American Geriatrics Society updated Beers Criteria for potentially inappropriate medication use in older adults. *Journal of the American Geriatrics Society*. 2012;60(4):616-631.
196. Lin JS, O'Connor E, Rossom RC, Perdue LA, Eckstrom E. Screening for Cognitive Impairment in Older Adults: A Systematic Review for the U.S. Preventive Services Task Force. *Annals of internal medicine*. 2013;159(9):601-612.
197. Michael YL, Whitlock EP, Lin JS, et al. Primary care-relevant interventions to prevent falling in older adults: a systematic evidence review for the U.S. Preventive Services Task Force. *Annals of internal medicine*. 2010;153(12):815-825.
198. Sullivan DR, Iyer AS, Enguidanos S, et al. Palliative Care Early in the Care Continuum among Patients with Serious Respiratory Illness: An Official ATS/AAHPM/HPNA/SWHPN Policy Statement. *American journal of respiratory and critical care medicine*. 2022;206(6):e44-e69.
199. Li NC, Lee A, Whitmer RA, et al. Use of angiotensin receptor blockers and risk of dementia in a predominantly male population: prospective cohort analysis. *BMJ (Clinical research ed)*. 2010;340:b5465.
200. Whitmer RA. Type 2 diabetes and risk of cognitive impairment and dementia. *Current neurology and neuroscience reports*. 2007;7(5):373-380.
201. Claytor JD, Grant RW. Reassessment of Glycemic Goals in Older Patients With Dementia-More Work Is Needed to Do Less. *JAMA internal medicine*. 2021;181(7):994-995.
202. Bayliss EA, Edwards AE, Steiner JF, Main DS. Processes of care desired by elderly patients with multimorbidities. *Family practice*. 2008;25(4):287-293.
203. Bayliss EA, Ellis JL, Shoup JA, Zeng C, McQuillan DB, Steiner JF. Effect of continuity of care on hospital utilization for seniors with multiple medical conditions in an integrated health care system. *Annals of family medicine*. 2015;13(2):123-129.
204. Portz JD, Bayliss EA, Bull S, et al. Using the Technology Acceptance Model to Explore User Experience, Intent to Use, and Use Behavior of a Patient Portal Among Older Adults With Multiple Chronic Conditions: Descriptive Qualitative Study. *Journal of medical Internet research*. 2019;21(4):e11604.
205. Tuzzio L, Berry AL, Gleason K, et al. Aligning care with the personal values of patients with complex care needs. *Health services research*. 2021;56 Suppl 1(Suppl 1):1037-1044.
206. National Institute on Aging. US Deprescribing Research Network. 2020; <https://deprescribingresearch.org/>. Accessed August 3, 2020.
207. Saadi A, Sanchez Molina U, Franco-Vasquez A, Inkelas M, Ryan GW. Assessment of Perspectives on Health Care System Efforts to Mitigate Perceived Risks Among Immigrants in the United States: A Qualitative Study. *JAMA network open*. 2020;3(4):e203028.
208. Radder DLM, Nonnekes J, van Nimwegen M, et al. Recommendations for the Organization of Multidisciplinary Clinical Care Teams in Parkinson's Disease. *Journal of Parkinson's disease*. 2020;10(3):1087-1098.
209. Larson EB, Stroud C. Meeting the Challenge of Caring for Persons Living With Dementia and Their Care Partners and Caregivers: A Report From the National Academies of Sciences, Engineering, and Medicine. *JAMA*. 2021;325(18):1831-1832.